

## **Abundance estimate of Australian east coast humpback whales (Group E1) in 2005 using multi year photo-identification data and capture-recapture analysis.**

DAVID A. PATON<sup>1,2,3</sup>, LYNDON BROOKS<sup>1,4</sup>, DANIEL BURNS<sup>1,2,4</sup>, ERIC KNIEST<sup>2,5</sup>, PETER HARRISON<sup>1,2,4</sup> AND PETER BAVERSTOCK<sup>1,4</sup>

<sup>1</sup> Southern Cross University, PO Box 157 Lismore, NSW 2480, Australia

<sup>2</sup> South Pacific Whale Research Consortium, Cook Islands Whale Research, Box 3069 Avarua, Rarotonga, Cook Islands, South Pacific

<sup>3</sup> Blue Planet Marine, PO Box 919 Jamison Centre, 2614, ACT Australia.

<sup>4</sup> Southern Cross University Whale Research Centre, PO Box 157 Lismore, NSW 2480, Australia

<sup>5</sup> University of Newcastle, School of Engineering, University Drive, Callaghan, NSW 2308 Australia

### **Abstract**

*Vessel based surveys were conducted between 1999 and 2005 during the annual humpback whale northern migration off Byron Bay on the east coast of Australia. A multi-year photo-identification sampling and capture-recapture analysis was undertaken for this 7 year period. This analysis resulted in a population estimate for the number of humpback whales that migrated north past Byron Bay in 2005 of 7390 (95% CI: 4040-10739). This estimate is consistent with other capture-recapture and land based estimates for Group E(1) and supports the growing body of data that indicates a high rate of increase for this population of humpback whales.*

### **Introduction**

In the Southern Hemisphere, humpback whales (*Megaptera novaeangliae*) undertake an annual migration during the austral winter months from their high-latitude summer feeding areas in the Antarctic to their low-latitude winter tropical breeding areas (Chittleborough 1965; Dawbin 1966). This northern annual migration is not random in structure, with clear temporal segregation of different classes of whales being recorded from whaling data. The first animals to leave the feeding grounds on the annual migration are lactating females and yearlings, followed by immature whales of both sexes, mature males and resting females at the peak of the season, and finally pregnant females (Dawbin 1966, 1997).

Humpback whale populations in the South west Pacific Ocean are made up of two populations (E and F). Group E has now been divided into three subgroups known as E(1) those wintering off the Australian east coast, E(2) those wintering around New Caledonia, an E(3) those wintering around Tonga (IWC 2005; Olavarria et al. 2006).

Following intensive over-exploitation of Southern Hemisphere humpback whale populations last century, including illegal whaling operations in the Southern Ocean (Clapham et al. 2009), Group E collapsed in 1962 to less than 5% of the original estimated pre-exploitation abundance (potentially in range of 30,000 to 40,000 humpback whales (Jackson et al. 2006)). Ongoing research over the last 27 years, monitoring the abundance of this humpback whale population has measured a high but steady rate of increase of 10.9% per annum (95% CI 10.5 – 11.4%) (Noad et al.

2008). This rate of increase is close to or at the estimated maximum biological rate of increase for this species (Clapham & Zerbini 2006). Other breeding stocks in the south west Pacific Ocean are showing limited or no signs of recovery (Gibbs and Childerhouse 2004; Garrigue et al., 2000, 2002; Paton et al. 2009; Paton et al. in review).

Recent estimates of adult survival rate for humpback whales include figures of 0.95 for the Gulf of Maine population (Buckland 1990, Clapham et al. 2003) and 0.96 for populations in the North Pacific Ocean (Calambokidis & Barlow 2004, Mizroch et al. 2004).

Research activities to monitor the abundance of the Group E(1) humpback whales have primarily relied on land based counts and distance sampling methodology (Bryden et al. 1988, 1996; Paterson et al. 1989, 1994, 2001, 2004; Brown, 1996, Brown et al. 2003; Noad et al. 2004, 2008). Noad et al. (2008) estimated absolute abundance of this stock based on land based counts for 2007 at 9,683 whales (95% CI: 8,556 – 10,959). Paton et al. (in press) undertook a single-season, multi-point sampling and capture–recapture analysis using photo-identification data for the population of migrating whales in 2005. These data were analysed using a variety of closed population models with a model-averaged estimate of 7,041 (95% CI: 4075 – 10,008) whales migrating along the east Australian coast in 2005. Although each of these abundance estimates relies on different assumptions and is subject to different inherent biases, both methods produced a similar population estimate (taking into consideration the reported annual population recovery rate of 10.9% (Noad et al. 2008)).

By using a number of techniques to assess the current abundance of the Group E(1) population, a more robust population estimate can be obtained. Here, we present a further estimate for the abundance of the humpback whales migrating along the Australian east coast in 2005 using a multi-year sampling and capture-recapture analysis.

## **Methods**

### *Survey methods*

Vessel based photo-identification surveys were undertaken annually at Byron Bay, (Latitude 28° 37'S, Longitude 153° 38'E) between 1999 and 2005. Byron Bay, located in northern New South Wales, is the most easterly point on the Australian mainland and is located on the edge of the migratory corridor of humpback whale population E1 (Figure 1). At this location the bulk of the migrating humpback whales migrate within 10km of the coast (Bryden 1985). The width of the humpback whale migration corridor was re-assessed in 1991 (Brown 1996) and 2007 (Noad and Dunlop 2007) at Point Lookout (135km north of Byron Bay), and found to be consistent with the results of Bryden (1985). Humpback whales travel past the eastern point of Australia at Byron Bay, in both a northerly and southerly direction, en-route between the Antarctic feeding grounds and the Great Barrier Reef breeding grounds (Paterson 1991).

Surveys were undertaken as whales migrated north past Cape Byron during the austral winter months (May – August). Survey effort was not consistent for each field season between 1999 and 2005, and while the annual surveys were timed to include what was

anticipated to be the peak of the northern migration passing Byron Bay, based on historical and recent data (Paterson 1991; Dawbin 1997; Paton and Kniest 2006), most annual surveys covered only a short part of the migration period.

The research vessel was assisted in finding pods of whales by a team of land-based observers working from Cape Byron lighthouse using the “Cyclopes” (theodolite/computer) whale tracking system (Kniest and Paton 2001). Survey effort, timing and equipment used during each annual survey are summarised in Table 1.

Photography of ventral fluke surfaces was obtained using a standard sampling protocol throughout the duration of the project. Ventral fluke photographs were obtained during a maximum of ten terminal dives and/or a maximum of 45 minutes with each pod (as per Smith et al. 1999). Photographs of the ventral fluke surfaces of calves were not included in this study.

#### *Photo-Identification*

The single best photograph for each individual whale for each field season was selected. Non digital images were scanned using a Nikon Coolscan III at maximum resolution. All images were cropped to a common 3 x 2 pixel ratio as high quality jpeg digital files. Composites of multiple images of a single fluke were constructed if these provided sufficient information to accurately identify the whale (Figure 2). All images for each field season were assessed for within-season resights to eliminate duplicates.

To address potential bias associated with photo quality, an independent cetacean researcher experienced in humpback whale fluke photograph matching, reviewed the fluke catalogue using a protocol developed in the Northern Hemisphere for grading humpback whale fluke identification photograph quality (Calambokidis *et al.* 2001). This included scoring all flukes according to five different characteristics of photo quality: (1) exposure / contrast / lighting; (2) fluke angle; (3) photographer / lateral angle; (4) focus / sharpness; and (5) proportion of fluke visible. Each photograph was given a score from 1 to 5 (highest quality to lowest) for each characteristic, and all flukes with at least one score of 4 or lower (5) were excluded from the dataset.

#### *Matching*

A stratified matching system developed by one of the authors (DB) was used in this analysis. This system is based on stratifying flukes by different characteristics including percentage black, characteristics of the centre and characteristics of the trailing edge of the fluke for each identification photograph. This system was used to reduce the number of comparisons required in the matching process. All matches found were reviewed and confirmed by at least two of the authors (DP and DB).

#### *Statistical model*

Only the 2005 sample, which extended over a large part of the total migration (a 10 week survey period), might be considered to be a random sample from the migrating population. The samples in the years 1999 to 2004 would have been selected from the classes of whales passing Byron Bay in the relatively short periods of the surveys in those years. This is unlikely to have been the same part of the population each year because a) the peak of the migration can change by up to five weeks from year to year

(Dawbin 1956), and b) some whales are likely to have changed age class or pregnancy status between years, and therefore the timing of their migration.

The series of surveys describes an open population with births, mortality, potential immigration and emigration occurring between years. However, no open population model was available that was appropriate for the data analysis.

We used the Chapman small sample modification to the Lincoln-Petersen two sample estimator for  $N$  (see e.g., Chao and Huggins 2005) based on a sample pooled over the years 1999 to 2004, and compared with the 2005 sample. This is a consistent estimator of the population size when either or both of the samples are a simple random sample as we assumed for the 2005 sample (see e.g., Schwarz 2005). We adjusted the number marked in the pooled sample ( $n_1$ ) to the number expected to have been alive and available for resighting in 2005 ( $n_{1A}$ ) by reducing the count from the year last sighted assuming a true survival rate of 0.95 pa. This procedure adjusted for the mortality between marking and recapture. Births and permanent immigrants in the interval 1999 to 2005 would have been represented in the 2005 sample. There was no basis on which to make any adjustment for permanent emigrants.

The Chapman estimator of population size ( $\hat{N}$ ):

$$\hat{N} = \frac{(n_1 + 1)(n_2 + 1)}{(m_2 + 1)} - 1$$

$$Var(\hat{N}) = \frac{(n_1 + 1)(n_2 + 1)(n_1 - m_2)(n_2 - m_2)}{(m_2 + 1)^2(m_2 + 2)}$$

where,

$n_1$  = number identified in first sample,

$n_2$  = number identified in second sample,

$m_2$  = number common to both samples.

## Results

Table 1 describes the survey dates for the years 1999 to 2005. As may be seen there only the 2005 survey extended over a relative large part of approximately 12 weeks of the migration period.

A total of 1008 fluke photographs were assessed for inclusion in the analysis. Following independent assessment against photograph quality protocols, 243 fluke photographs were excluded from the data set based on photographic quality. The final data set comprised a total of 765 fluke identification photographs. Table 2 reports the matches and frequencies of capture histories. Table 3 reports the number of whales sighted between 1999 and 2004 ( $n_1$ ) and the number of those expected to have been alive and available for capture in 2005 ( $n_{1A}$ ) based on an assumed mortality rate of 5%.

Three hundred and thirty seven (337) whales were sighted in 2005 ( $n_2$ ) of which 15 had been previously sighted at least once in the period 1999-2004 ( $m_2$ ). The Chapman

estimate of the population size in 2005 ( $\hat{N} | n_1, n_2, m_2$ ) was 8,850 (95% CI: 4823-12878) whales.

The mortality-adjusted Chapman estimate of the population size in 2005 ( $\hat{N} | n_{1A}, n_2, m_2$ ) was 7390 (95% CI: 4040-10739) whales.

## Discussion

This study represents a multi-year sampling and capture-recapture abundance estimate using photo-identification for Group E(1) humpback whales migrating north past Byron Bay on the east coast of Australia during 2005. This estimate of 7,390 (95% CI: 4040-10739) humpback whales is consistent with the Paton et al. (in press) multi-point sampling and capture-recapture estimate for 2005 of 7,041 (95% CI: 4075 - 10,008) and Noad et al. (2008) land based distance sampling estimate of 9,683 whales (95% CI: 8,556:10,959) for 2007 for this same population. The Noad et al. (2008) estimate, when extrapolated backwards using an annual increase rate of 10.9%, would give an estimate for 2005 of 7,873 (95% CI: 6,597-8,911).

## Acknowledgements

This study was undertaken with support from New South Wales National Parks and Wildlife Service, Southern Cross University, Southern Cross University Whale Research Centre, University of Newcastle, Cape Byron Headland Reserve Trust and Blue Planet Marine. This research was conducted under Scientific Research Permits issued by the Department of the Environment, Water, Arts and Heritage under the EPBC Act 1999 (permit number E2001/0005) and the New South Wales National Parks and Wildlife Service (permit number S10403).

The authors would like to thank all the volunteers who assisted in the Cape Byron Whale Research Project with special thanks to Adrian Oosterman and Merv Whicker. Our thanks also go to Rochelle Constantine for undertaking the quality control assessment of the fluke identification catalogue and Greg Gorman for assisting with fluke matching. The authors would like to Greg Luker for providing Figure 1.

## Literature Cited

Blackmer A.L., Anderson S.K. and Weinrich M.T. (2000) Temporal variability in features used to photo-identify humpback whales (*Megaptera novaeangliae*). *Marine Mammal Science* 16(2): 338-354

Brown, M.R. 1996. Population biology and migratory characteristics of east Australian Humpback whales, *Megaptera novaeangliae*. PhD thesis University of Sydney. (unpublished). pp 1-322. [Available from the University of Sydney].

Brown, M. R., Corkeron, P. J. Hale P.T., Schultz, K.W. and Bryden M.M. 1995. Evidence for a sex segregated migration in the humpback whale (*Megaptera novaeangliae*). *Proceedings of the Royal Society of London*. B 259: 229-234.

Brown M.R., Field, M.S. Brown, C.E and Bryden, M.M. 2003. Rates of increase for east Australian humpback whales, *Megaptera novaeangliae*, form 1991 -2000. Paper SC/55/SH21 presented to the IWC Scientific Committee, 1989 (unpublished).

Bryden , M. M. 1985. *Studies of humpback whales (Megaptera novaeangliae), Area V. Studies of sea mammals in south latitudes*. Proceeding of the 52nd ANZAAS Congress. J. K. Ling and M. M. Bryden. Sydney, Australia: 115-124.

- Buckland, S. T. 1990. Estimation of survival rates from sightings of individually identifiable whales. *Report of the International Whaling Commission* (Special Issue 12):149–153
- Calambokidis, J. and Barlow, J. 2004. Abundance of blue and humpback whales in the eastern North Pacific estimated by capture-recapture and line-transect methods. *Marine Mammal Science* 20(1): 63–85.
- Calambokidis, J., Steiger, G. H., Straley, J. M., Herman, L. M., Cerchio, S., Salden, S. R., Urbán, J., Jacobsen, J. K., von Zeigesar, O., Balcomb, K. C., Gabriele, C. M., Dahlheim, M. E., Uchida, S., Ellis, G., Miyamura, Y., de Guevara, P. L., Yamaguchi, M., Sato, F., Mizroch, S. A., Schlender, L., Rasmussen, K., Barlow, J., and Quinn II, T. J. 2001. Movements and population structure of humpback whales in the North Pacific. *Marine Mammal Science* 17 (4), 769-794.
- Carlson C.A., Mayo C.A. and Whitehead W. (1990) Changes in the ventral fluke pattern of the humpback whale (*Megaptera novaeangliae*), and its effects on matching; Evaluation of its significance to photo-identification research. *Report of the International Whaling Commission* Special Issue 12: 105-111.
- Chittleborough, R. G. 1965. Dynamics of two populations of the humpback whale, *Megaptera novaeangliae* (Borowski). *Australian Journal of Marine and Freshwater Research*. 16: 33-128.
- Clapham, P., Mikhalev, Y., Franklin, W., Paton, D.A., Baker, C.S., Ivashchenko, Y. and Brownell R.L. (2009) Catches of humpback whales, *Megaptera novaeangliae*, by the Soviet Union and other Nations in the Southern Ocean, 1947-1973. *Marine Fisheries Review* 71(1):39-43
- Clapham, P., and Zerbini, A. 2006. Is social aggregation driving high rates of increase in some Southern Hemisphere humpback whale populations? Paper SC/58/SH3 presented to the IWC Scientific Committee, 2006 (unpublished). 11pp.
- Clapham, P. J., J. Barlow, M. Bessinger, T. Cole, D. Mattila, R. Pace, D. Palka, J. Robbins and R. Seton. 2003. Abundance and demographic parameters of humpback whales from the Gulf of Maine, and stock definition relative to the Scotian Shelf. *Journal of Cetacean Research and Management* 5:13–22.
- Chao, A. and Huggins R. 2005. Classical closed-population capture-recapture models. In Steven C. Amstrup, Trent L. McDonald, and Bryan F.J. Manly (Eds) *Handbook of Capture-Recapture Analysis*. Princeton University Press, NJ.
- Dawbin, W. H. 1956. The migrations of humpback whales which pass the New Zealand coast. *Transactions and Proceedings of the Royal Society of New Zealand*. 84: 147-96.
- Dawbin, W. H. 1966. The seasonal migratory cycle of humpback whales. pp. 145-171. In: K.S. Norris (ed.) *Whales, dolphins and porpoises*. University of California Press, Berkeley:.
- Dawbin, W. H. 1997. Temporal segregation of humpback whales during migration in Southern Hemisphere waters. *Memoirs of the Queensland Museum* 42: 105.
- Garrigue, C., A. Aguayo, Amante Helweg, V.L.U., Baker, C.S., Caballero, S., Clapham, P., Constantine, R., Denking, J., Donoghue, M., Florez-Gonzalez, L., Greaves, J., Hauser, N., Olavarria, C., Pairoa, C., Peckham, H. and Poole, M. 2002. Movements of humpback whales in Oceania, South Pacific. *Journal of Cetacean Research and Management*. 4: 255-260.
- Garrigue, C., Forestell, P., Greaves, J., Gill, P., Naessig, P., Patenaude, N.M. and Baker, C.S. 2000. Migratory movements of humpback whales (*Megaptera novaeangliae*) between New Caledonia, east Australia and New Zealand. *Journal of Cetacean Research and Management*. 2: 111-115.
- Gibbs N. and Childerhouse S. 2004. First report of the recovery of humpback whales in New Zealand waters. (unpublished) [Available from the Department of Conservation. PO Box 10 420 Wellington, New Zealand]

- International Whaling Commission. 2005. Inter-sessional Working Group on Southern Hemisphere Humpback Whales: Revised Table by Breeding Stock (as at 1 May 2005). Paper SC/57/SH11 presented to the IWC Scientific Committee, May 2006 (unpublished).
- Jackson, A.J., Zerbini, A., Clapham, P. Garrigue, C., Hauser, N., Poole, M. and Baker, C.S. 2006. A Bayesian assessment of humpback whales on breeding grounds of Eastern Australia and Oceania (IWC Stocks E, E1, E2 and F). Paper SC/A06/HW52 presented to IWC Scientific Committee, May 2006 (unpublished).
- Kniest, E. and Paton, D.A. 2001. Temporal GIS for Marine Mammal Research. (Abstract). 14 Biennial Conference on the Biology of Marine Mammals. Vancouver, Canada. (Society for Marine Mammalogy)
- Mizroch, S. A., Herman, L.M., Straley, J.M., Glockner-Ferrari, D.A., Jurasz, C., Darling, J., Cerchio, S., Gabriele, C.M., Salden, D.R., von Ziegesar, O., 2004. Estimating the adult survival rate of central North Pacific humpback whales (*Megaptera novaeangliae*). *Journal Of Mammalogy* 85(5): 963-972.
- Noad, M. and Dunlop, R. 2007. Abundance estimates of the east Australian humpback whale population. Progress (unpublished) Report for the Australian Department of the Environment and Water Resources. [Report available from Australian Department of the Environment and Water, Arts and Heritage, GPO Box 787 Canberra, 2601 ACT Australia].
- Noad, M.J., Paton, D.A and Cato, D.E. 2005. Absolute and relative abundance estimates of Australian east coast humpback whale (*Megaptera novaeangliae*). Paper SC/A06/HW27 presented to the IWC Scientific Committee, 2005 (unpublished).
- Noad, M.J., Paton, D.A., Cato, D.H, Dunlop, R., Kniest, E. and Wyn Morris, C. 2006. Survey of East Australian Humpback Whales (*Megaptera novaeangliae*) from Point Lookout 2004. Unpublished report to the Department of Environment and Heritage, Canberra. Australia. [Report available from Australian Department of the Environment and Water, Arts and Heritage, GPO Box 787 Canberra, 2601 ACT Australia].
- Noad, M.J., Dunlop, R., Paton, D. and Cato, D.H. (2008) An update of the east Australian humpback whale population (E1) rate of increase. SC/60/SH31
- Olavarria, C., Anderson, M., Paton, D.A., Burns, D., Brasseur, M., Garrigue, C., Hauser, N., Poole, M., Caballero, S., Florex-Gonzalez, L. and Baker, C.S. 2006. Eastern Australia humpback whale genetic diversity and their relationship with Breeding Stocks D, E, F and G. Paper SC/58/SH25 presented to the IWC Scientific Committee, June 2006 (unpublished) 6pp presented to the IWC Scientific Committee, 2005 (unpublished).
- Paterson, R. A. 1991. The migration of Humpback Whales *Megaptera novaeangliae* in east Australian waters. *Memoirs of the Queensland Museum* 30(2):333-341.
- Paterson, R. A., P. Paterson and Cato, D.H. 2001. Status of humpback whales, *Megaptera novaeangliae*, in east Australia at the end of the 20th century. *Memoirs of the Queensland Museum* 47(2):579-586.
- Paton, D., Batibasaga, A., Shama, S., O'Connor, W. and Nand, N. (2009). Report of the Whale and Dolphin Survey undertaken in the Lomaiviti Island Group, Fiji 2008 [Available from Blue Planet Marine, PO Box 919 Jamison Centre, ACT 2614 Australia] (unpublished).
- Paton, D.A., Brooks, L., Burns, D., Franklin, T., Franklin, W., Harrison, P. and Baverstock, P. (in press) Abundance of east coast Australian humpback whales (*Megaptera novaeangliae*) in 2005 estimated using multi-point sampling and capture-recapture analysis. *Journal of Cetacean Research and Management*.
- Paton, D.A. and Kniest, E. (in review) Population growth of Australian east coast humpback whales, observed from Cape Byron, 1998 to 2004. Submitted to *Journal of Cetacean Research and Management*.

Paton, D.A., Oosterman, A., Whicker, M., Kenny I., Christian, M. and Garrigue, C. (In review) Assessment of sighting survey data for humpback whales (*Megaptera novaeangliae*) at Norfolk Island 2003-2006 and a comparison with historical records for the region. Submitted to *Journal of Cetacean Research and Management*.

Rice, M., Carlson, C., Chu, K., Dolphin, W. and Whitehead, H. 1987. Are humpback whale population estimates being biased by sexual differences in fluking behavior? Report of the Scientific Committee. *International Whaling Commission*. 37: 333-335.

Carl J. Schwarz. 2005. Multistate models. In Steven C. Amstrup, Trent L. McDonald, and Bryan F.J. Manly (Eds) *Handbook of Capture-Recapture Analysis*. Princeton University Press, NJ.

Smith, T. D., Allen, J., Clapham, P. J., Hammond, P. S., Katona, S., Larsen, F., Lien, J., Mattila, D., Palsboll, P. J., Sigurjonsson, J., Stevick, P. T. and Oien, N. 1999. An ocean-basin-wide mark-recapture study of the North Atlantic humpback whale (*Megaptera novaeangliae*). *Marine Mammal Science*. 15(1):1-32.

**Table 1 Summary of survey effort and equipment used.**

Year	1999	2000	2001	2002	2003	2004	2005
Dates of survey	19- 20 June, 30 June – 11 July	25 June – 9 July	23 June – 8 July	17 June – 20 July	21 June – 6 July	26 June – 11 July	4 June – 12 Aug
Survey period (days)	14	14	16	34	16	16	69
Research vessel	5.8 metre centre console	5.8 metre centre console	5.8 metre centre console	5.8 metre centre console	5.8 metre centre console	5.8 metre centre console	5.5 metre centre console
Photographic equipment	Canon EOS 1, 300mm lens F2.8	Canon EOS 1, 300mm lens F2.8	Canon EOS 10D, 100-400mm lens F3.5-5.6IS	Canon EOS 10D, 100-400mm lens F3.5-5.6IS	Canon EOS 10D, 100-400mm lens F3.5-5.6IS	Nikon D100, 70-200mm lens F2.8 VR and 1.4x converter	Canon EOS 20D, 100-400mm lens F3.5-5.6IS

**Table 2 Matches and frequencies for capture histories of humpback whales recorded near Byron Bay, Australia.**


Row	1999	2000	2001	2002	2003	2004	2005	n_row
1	1	0	0	0	0	0	0	32
2	0	1	0	0	0	0	0	32
3	0	0	1	0	0	0	0	64
4	0	0	0	1	0	0	0	103
5	0	0	0	0	1	0	0	61
6	0	0	0	0	0	1	0	101
7	0	0	0	0	0	0	1	322
8	1	0	0	1	0	0	0	1
9	1	0	0	0	0	1	0	1
10	1	0	0	0	0	0	1	1
11	0	1	0	1	0	0	0	1
12	0	0	1	1	0	0	0	1
13	0	0	1	0	1	0	0	1
14	0	0	0	1	1	0	0	1
15	0	0	0	1	0	1	0	1
16	0	0	0	1	0	0	1	7
17	0	0	0	0	1	1	0	3
18	0	0	0	0	1	0	1	2
19	0	0	0	0	0	1	1	5

**Table 3** Number of sampled humpback whales estimated to be alive in 2005 based on an estimated 0.95 population survival rate.

Year	n-last sighted	Estimated n-alive in 2005 <sup>1</sup>
1999	32	23.52
2000	32	24.76
2001	64	52.13
2002	106	90.88
2003	63	56.86
2004	106	100.70
1999-2004	$n_1 = 403$	$n_{1A} = 348.85$

<sup>1</sup> Calculated as  $(n\text{-last sighted}) \times (0.95)^{(2005\text{-Year})}$

**Figure 1** Study Area



**Figure 2.** Example of a composite image used in the analysis.

