

GENETIC DIVERSITY AND POPULATION STRUCTURE OF HUMPBACK WHALES (*Megaptera novaeangliae*) FROM ECUADOR BASED ON MITOCHONDRIAL DNA ANALYSES. UPDATE OF SC/59/SH11

FERNANDO FÉLIX¹, SUSANA CABALLERO^{2,3} AND CARLOS OLAVARRÍA^{2,4}

¹Fundación Ecuatoriana para el Estudio de Mamíferos Marinos (FEMM). PO Box 09-01-11905, Guayaquil, Ecuador.

²School of Biological Sciences, University of Auckland, Auckland, New Zealand.

³Present address: Universidad Los Andes, Bogotá, Colombia.

⁴Fundación Centro de Estudios del Cuaternario Fuego-Patagonia y Antártica (CEQUA), c/o INACH, Plaza Muñoz Gamero 1055, Punta Arenas, Chile.

E-mail: fernandofelix@femm.org

ABSTRACT

Information on the genetic characterization of humpback whales (*Megaptera novaeangliae*) breeding off Ecuador (2°10'S, 81°00'W; Breeding Stock G) is presented. Mitochondrial DNA was extracted and sequenced from 103 skin samples collected between 2002 and 2007 to establish the genetic diversity of Ecuadorian humpback whales. Samples were obtained either from beached animals ($n=4$), biopsies ($n=1$) or sloughed skin ($n=98$). Twenty nine different haplotypes were found, five of which were new and unique. Haplotype diversity ($h \pm SD$) was estimated to be 0.893 ± 0.023 and the nucleotide diversity ($\pi \pm SD$) 0.018 ± 0.009 . A pair-wise analysis of molecular variance (AMOVA) was used to compare diversity within and between other areas of known distribution for this stock in the Southeast Pacific (Colombia and Magellan Strait) and the Antarctic Peninsula. Significant differentiation at both haplotype and nucleotide levels was found only with Magellan Strait ($p < 0.0001$). When data from 2006 and 2007 were stratified by sex and year, significant differences were found at both haplotype and nucleotide level between females in 2006 and females in 2007 ($p < 0.01$) and between females in 2006 and males in 2007 ($p < 0.05$). Although the pooled dataset analysis suggests panmixia in the Breeding Stock G, stratified data by sex and year indicate some level of heterogeneity, possibly due to differential female migrating behaviour. Such heterogeneity would be also responsible of the skewed sex proportion toward males obtained (2.5:1). Further research and a larger number of samples from different sites are required to assess appropriately the structure of this population.

KEY WORDS: humpback whale, genetics, breeding grounds, South America, Breeding Stock G.

INTRODUCTION

Humpback whales (*Megaptera novaeangliae*) distribute during the austral winter along the northwestern coast of South America, mainly off Colombia and Ecuador, but also further north, off Panama (Kellog, 1929; Townsend, 1935; Mackintosh, 1942). Whales wintering off those neighbouring locations were considered to form a single breeding stock, despite the lack of information supporting that assumption. Only in the last decade the putative single breeding stock hypothesis was addressed, when individually photo-identified whales were compared among those areas (Flórez-González *et al.*, 1998). Despite the reduced number of compared individuals from Ecuador, a few whales were matched, which suggested that interchange among these regions occurred. Recently, humpback whales have been recorded during the austral winter further north, off Costa Rica (Acevedo-Gutiérrez and Smultea, 1995;

Rasmussen *et al.*, 2007), which expands the range of the wintering grounds of this species in the Southeast Pacific population. All those breeding areas have been linked to the feeding areas in the west side of the Antarctic Peninsula (Stone *et al.*, 1990; Caballero *et al.*, 2001; Garrigue *et al.*, 2002; Stevick *et al.*, 2004; Rasmussen *et al.*, 2007) and the Magellan Strait, in southern Chile (Acevedo *et al.*, 2007; Capella *et al.* 2008).

The discreteness of Southeast Pacific humpback whale population among the Southern Hemisphere metapopulation was also assumed for a long time (Kellog, 1929; Mackintosh, 1942; Omura, 1953), despite, again, any evidence to support this. Only recently, based on photo-identification (Garrigue *et al.*, 2002; Stevick *et al.*, 2004) and genetic analyses (Caballero *et al.*, 2001; Olavarria *et al.*, 2007), the putative discreteness of this breeding stock (Breeding Stock G; IWC, 1998) was confirmed by comparisons with neighbouring Southern Hemisphere breeding stocks.

Genetic studies on the species have been conducted in recent years in different locations in the Southeast Pacific, including breeding grounds off mainland Ecuador and the Galapagos Islands (Félix *et al.*, 2006a, 2007a), Gorgona Island and Málaga Bay in Colombia (Caballero *et al.*, 2000, 2001; Olavarria *et al.*, 2007) and feeding areas at the Magellan Strait in southern Chile (Olavarria *et al.*, 2006), and along the west coast of the Antarctic Peninsula (Olavarria *et al.*, 2000). Such studies have provided an overview of the genetic diversity, which seems to be the lowest among humpback whale stocks in the Southern Hemisphere (Olavarria *et al.*, 2007), and have shown a lack of genetic differentiation between the Antarctic Peninsula feeding area and the Colombian breeding ground (Caballero *et al.*, 2001; Olavarria *et al.*, 2000; 2007). Interestingly, the whales inhabiting the Magellan Strait, represent a separate feeding aggregation (Acevedo *et al.*, 2007), with their own genetic distinctiveness (Olavarria *et al.*, 2006).

Despite these recent contributions, there are gaps on our knowledge of this humpback whale stock, particularly about its population structure. Both genetics and photo identification studies (Olavarria *et al.*, 2006; Acevedo *et al.*, 2007) suggest some level of heterogeneity in the Breeding Stock G, comparable to what occurs in other Southern Stocks (see Rosenbaum *et al.*, 2006). Sightings of humpback whales almost year round off Peru (Ramírez, 1988) also suggest that not every animal from this stock complete the annual migration; some animals may stay south of the breeding area, or north of the feeding area if the case, in the high productive waters of the Humboldt Current off Peru and Chile, where they could find food in a predictable manner (Papastavrou and Van Waerebeek, 1997). This migrating behavior would not be exclusive from this stock since it has been reported elsewhere (e.g. Craig and Herman, 1997; Papastavrou and Van Waerebeek, 1997)

In this report we expand our previous analyses on the mitochondrial DNA (mtDNA) control region genetic diversity of humpback whales from Ecuador (see Félix *et al.*, 2006a, 2007a). Our findings show differences at haplotype and nucleotide composition showing a higher level of structure in this population than previously known. We also confirm a lack of differentiation among Ecuador and Colombia breeding grounds and the Antarctic Peninsula feeding area, but significant differences with the Magellan Strait feeding area.

MATERIALS AND METHODS

Sampling

Humpback whale skin samples were obtained between 2002 and 2007 off Ecuador. Four samples were collected from beached animals, 127 from sloughed skin (Amos *et al.*, 1992) and one from biopsing with a Barnett crossbow equipped with a 60cm long arrow and modified tip (Lambertsen, 1987). Three beached animals from mainland Ecuador and the

biopsied whale from Galapagos Islands were reported previously (Félix *et al.*, 2006b). Sloughed skin samples were obtained during the breeding seasons 2006-2007 (July-October) onboard whalewatching vessels working at Salinas, Ecuador (2°10'S, 81°00'W) (Figure 1). Sampling was conducted by a research team as part of a long-term research program by the Ecuadorian Foundation for the Study of Marine Mammals – FEMM (see Félix and Haase, 2001, 2005 for additional reference on the study).

When sampling for sloughed skin, the whale-watching boat skippers were asked to approach the site where a whale entered the water after an energetic surface display. Then small pieces of skin were picked up from the upper water column with a net attached to a pole. Pieces of skin were retrieved from the net and stored in plastic containers with either a solution of DMSO saturated NaCl or ethanol 50%. The net was thoroughly washed with sea water until no pieces of skin were visible on its surface, and then the device was considered ready for the next sampling attempt. Once on shore, samples were stored at 4°C for up to six months prior to laboratory analysis.

Sampled whales were photographed for individual identification with the pigmentation pattern on the ventral side of flukes (photo-identification, Katona *et al.*, 1979). It was possible to photo-identify, however, only 65% of the sampled whales.

Usually only one animal was sampled by group but occasionally it was possible to collect two or three samples, presumably from different individuals. When more than one sample was taken from the same group the results of the genetic analyses were necessary to define its distinctiveness. If it resulted in the same sex and haplotype, then duplicity was presumed and the sample was not included in the statistical analysis. This criterion was not applied to a mother with female calf. The bias introduced by duplicity would be in the same order of the within year resighting rate obtained simultaneously with photo-ID, this was 6.4% in the season 2006 (Félix *et al.*, 2007b). However, from photo-ID records it was established that only one whale was sampled twice (1.5% of the identified whales).




Figure 1. The study area on the coast of Ecuador.

Molecular analyses

A fragment of approximate 800bp of the mitochondrial DNA control region (CR) was amplified via the Polymerase Chain Reaction (Saiki *et al.*, 1988) using standard reaction conditions (Palumbi, 1996). For the PCR, we used the primer combination t-Pro-whale Dlp1.5 (5'-TCACCCAAAGCTGRARTTCTA-3') and Dlp8 (5'-CCATCGWGATGTCTTATTTAAGRGGAA-3') (Baker *et al.*, 1998; Olavarría *et al.*, 2007). The PCR profile was as follows: an initial denaturation at 95°C for 2 minutes, 36 cycles of 94°C for 30 seconds, 55°C for 1 minute and 72°C for 1 minute and 30 seconds, and a final extension at 72°C for 5 minutes. Free nucleotides and primers were removed from the PCR products using the PCR Cleaning kit (Invitrogen). PCR products were sequenced in both directions using the standard protocols of Big Dye™ terminator sequencing chemistry on an ABI 3100 automated capillary sequencer (Perkin Elmer), using the same PCR primers.

All sequences were manually edited and aligned using Sequencer 4.1 software (Gene Codes Corporation). Sequences were trimmed to 469bp to match a consensus region analysed previously (Olavarría *et al.*, 2006; 2007). Control region sequences were compared using MacClade (Maddison and Maddison, 2000) to identify haplotypes. Ecuador haplotypes were compared with haplotypes previously identified in other humpback whale populations in the South Pacific (Olavarría *et al.*, 2006, 2007).

Specific sex markers for gender determination followed the methodology of Gilson *et al.* (1998), which amplify a fragment (224 bp) of the *SRY* gene located on the Y chromosome. As internal positive control against PCR amplification failure, homologous ZFY/ZFX region (445bp) were amplified. Thus, in the electrophoresis analysis two bands of 224 and 445 bp were be present in males and only one in females (445bp).

Data analyses

Genetic diversity at haplotype (H) and nucleotide (π) levels were computed with the software Arlequin Ver 3.1 (Schneither *et al.*, 2006). Within population differentiation in haplotype frequencies (F_{ST}) and nucleotide (Φ_{ST}) composition as well as at geographic level between Ecuador and Colombia, Antarctic Peninsula and Magellan Strait was quantified using an Analysis of Molecular Variance (AMOVA) (Excoffier, 1995). Stratified data from 2006 and 2007 by sex and year were additionally tested with an exact test of sample differentiation based on haplotype frequencies (Raymond and Rousset, 1995). Both AMOVA and exact test are implemented in the Arlequin software.

RESULTS

Genetic diversity

From the 132 samples obtained, 14 failed sequencing and 15 were considered as duplicates. Thus 103 samples were considered in subsequent analyses, which revealed twenty-nine haplotypes (Table 1). The variable nucleotides include two insertion/deletions, 32 transitions and 1 transversion. Haplotype diversity ($h \pm SD$) was estimated to be 0.893 ± 0.023 and the nucleotide diversity (π) 0.018 ± 0.009 . The mean of pair-wise differences was 8.44 ± 3.93 .

When compared with a total of 120 haplotypes from other Southern Hemisphere breeding grounds and feeding areas (Olavarría *et al.*, 2006, 2007), 24 Ecuador haplotypes matched. The remaining five haplotypes were new and corresponded to single Ecuadorian individual whales (Table 2). Five of the previously reported haplotypes (SP16, SP19, SP26, SP41 and SP42) had been not found in either a breeding ground or a feeding area of the Stock G.

Interestingly SP16 haplotype had been only reported in the Western Australian population (see Olavarria *et al.*, 2007).

Table 1. Variable sites of 5 new haplotypes from Ecuador when compared with 120 humpback whales mtDNA control region sequences (Olavarria *et al.*, 2006, 2007). Alignment in relation to SP1 haplotype.

Haplotype	Variable sites											
	1	1	1	2	2	2	2	2	3	3		
	2	6	8	9	9	3	6	6	4	6	6	7
SP1	G	T	C	T	A	A	T	C	C	G	C	T
Mno25Ec06	.	C	T	.	C	G	.	T	T	.	T	.
Mno32Ec06	C
Mno35Ec06	C	G	.	.	T	A	.	C
Mno52Ec06	.	C	T	.	C	G	.	.	T	.	T	C
Mno37Ec07	A	G	C	T

Table 2. Ecuador humpback whale haplotype diversity and frequency of mtDNA control region sequences and proportion of haplotypes by sex and year (2006 and 2007). Haplotype nomenclature follows Olavarria *et al.* (2006, 2007).

Haplotype (469 pb)	Females				Males				Unknown sex	Overall	
	2006	2007	Total	%	2006	2007	Total	%		n	%
SP1		1	1	4,0	4	1	5	7,6		6	5,8
SP10						1	1	1,5		1	1,0
SP14					1		1	1,5		1	1,0
SP16	1		1	4,0						1	1,0
SP19						3	3	4,5		3	2,9
SP25	1		1	4,0	2	2	4	6,1	1	6	5,8
SP26	1		1	4,0						1	1,0
SP32		2	2	8,0	3	3	6	9,1	1	9	8,7
SP33	1		1	4,0	2		2	3,0		3	2,9
SP41		1	1	4,0						1	1,0
SP42	1		1	4,0	1		1	1,5		2	1,9
SP50		1	1	4,0	2		2	3,0		3	2,9
SP52					1		1	1,5		1	1,0
SP60	1		1	4,0	3	1	4	6,1		5	4,9
SP61	1		1	4,0	1		1	1,5	1	3	2,9
SP62					2	1	3	4,5		3	2,9
SP68		2	2	8,0		1	1	1,5		3	2,9
SP72						1	1	1,5		1	1,0
SP73	1		1	4,0		1	1	1,5		2	1,9
SP90		4	4	16,0	9	10	19	28,8	7	30	29,1
SP98		3	3	12,0	1	3	4	6,1	1	8	7,8
SP100						1	1	1,5		1	1,0
SP101						1	1	1,5		1	1,0
Mno03MA02						2	2	3,0	1	3	2,9
MnoEc2506					1		1	1,5		1	1,0
MnoEc3206					1		1	1,5		1	1,0
MnoEc3506	1		1	4,0						1	1,0
MnoEc3707		1	1	4,0						1	1,0
MnoEc5206	1		1	4,0						1	1,0
Total	10	15	25	100,0	34	32	66	100,0	12	103	100,0

Sex identification

The sex identification analysis revealed a significant sex bias towards males (68 males, 27 females; $\chi^2=17.7$, $p<0.01$). The same proportion was also found when only calves were considered (from seven sampled calves, five resulted males and two females).

Population structure by sex

A comparison of haplotype composition by sex was made in order to establish possible variability within the population. For this purpose information from 91 individuals with known haplotype and sex (25 females and 66 males) sampled in 2006 and 2007 was used. Through AMOVA tests the haplotype composition of females and males separately (two groups) as well as between sexes and years (four groups) were compared.

For females there were ten different haplotypes in 2006 and eight in 2007, whereas in males there were 15 different haplotypes in both years (Table 2). Most haplotypes were shared by both sexes and the proportion in most cases was similar, although the haplotype SP90 was almost 50% less represented in females. There was not a significant difference in haplotype frequency and nucleotide composition between sexes ($F_{ST} = -0.003$, $p = 0.535$ and $\Phi_{ST} = -0.0036$, $p = 0.578$).

When comparison included sex and year some differences arose, particularly in females. Haplotypes found in females were different between 2006 and 2007. It is surprising that the most common haplotype in the Breeding Stock G (SP90) was not present in females in 2006, although in the overall result this and other common haplotypes (e.g. SP32, SP25 and SP98) were well represented. In contrast, males showed a fairly regular haplotype frequency in both years. Significant differences in both haplotype frequency and nucleotide composition were found between females in 2006 and females in 2007 ($p < 0.01$ in both cases), as well as between females in 2006 and males in 2007 ($p < 0.05$ in both cases) (Table 3).

Table 3. Pair-wise test of differentiation for mtDNA control region sequence by sex and year based on the F_{ST} and Φ_{ST} indices. F = females, M = males, periods 2006 and 2007. The significance was analyzed using 5,000 non-parametric permutations of the data matrix.

	F 2006		F 2007		M 2006	
	F_{ST}	p -value	F_{ST}	p -value	F_{ST}	p -value
F 2007	0.1700	0.0083 \pm 0.0014				
M 2006	0.0758	0.0506 \pm 0.0030	-0.0072	0.4886 \pm 0.0075		
M 2007	0.1126	0.0157 \pm 0.0018	-0.0062	0.4650 \pm 0.0073	-0.0089	0.6175 \pm 0.006

	F 2006		F 2007		M 2006	
	Φ_{ST}	p -value	Φ_{ST}	p -value	Φ_{ST}	p -value
F 2007	1.6229	0.0085				
M 2006	0.6752	0.0591	-0.0329	0.4393		
M 2007	1.0756	0.0183	-0.0253	0.4165	-0.0749	0.6175

A global exact test of differentiation gave a p -value = 0.07960 \pm 0.0344 (20,000 Markov chain steps). However, when the differentiation test included samples by sex and year results were similar to those found based on the F_{ST} and Φ_{ST} indices (a highly significant difference between females in 2006 and females in 2007 ($p < 0.01$) and a significant difference between

females in 2006 and both males in 2006 and males in 2007 ($p < 0.05$); 100,000 Markov chain steps).

Population differentiation analysis with other Southeast Pacific areas

Ecuadorian haplotype frequencies were compared with other locations in the Southeast Pacific in both breeding (Colombia) and feeding areas (Magellan Strait and the Antarctic Peninsula) as reported by Olavarria *et al.*, (2006, 2007). The most common haplotype reported in Ecuador, SP90 (29.1%), is the most common haplotype in the Southeastern Pacific. The second and third most common haplotypes found in Ecuador SP32 (8.7%) and SP98 (7.8%) were also among the most common haplotypes in Colombia and Antarctic Peninsula, but were absent from Magellan Strait. The second most common haplotype from Magellan Strait was recorded in three whales from Ecuador, but was absent in Colombia and Antarctic Peninsula. Two relatively common haplotypes in Colombia SP52 y SP10 were recorded once in Ecuador; the former was found with similar frequency in Colombia and Antarctic Peninsula. Haplotype SP25 was recorded in 6 whales in Ecuador, but only once in Colombia and was absent in Antarctic Peninsula and Magellan Strait. The haplotype found in Galapagos (SP61), was recorded twice off mainland Ecuador, once off Colombia and two at Antarctic Peninsula. Overall, Ecuador humpback whales shared 17 haplotypes of 27 previously reported in Colombia (63%), 15 of 25 in Antarctic Peninsula (60%) and three of four from Magellan Strait (75%).

A pair-wise AMOVA between Ecuador and the other Stock G locations calculated a between variance of 5.97% and a within variance of 94.03%. The high proportion of the within variance indicates a high genetic similarity between the compared sites, as expected for a panmictic population. However, a significant difference was found between Ecuadorian and Magellan Strait whales in both haplotype frequency and nucleotide composition ($p < 0.001$ in both cases) (Table 4).

Table 4. Pair-wise test of differentiation for mtDNA control region sequence between Ecuador and other Stock G humpback whales based on the F_{ST} and Φ_{ST} indices. The significance was analyzed using 5,000 non-parametric permutations of the data matrix.

	Colombia	Magellan Strait	Antarctic Peninsula
F_{ST}	0.0023	0.1670	0.00206
p -value	0.1979	< 0.0001	0.2416
Φ_{ST}	0.00210	0.14464	0.0018
p -value	0.1966	<0.0001	0.2430

DISCUSSION

The analysis presented here is the first attempt to assess the putative panmixia within the breeding Stock G, including for first time in the analyses samples from Ecuador. Ecuadorian genetic diversity was similar to those from Colombia and Antarctic Peninsula areas as reported by Olavarria *et al.* (2006, 2007), supporting that the Stock G has the lowest genetic diversity among humpback whales in the South Pacific. In fact, the mtDNA diversity of the Breeding Stock G would be the lowest in the entire Southern Hemisphere with the exception the North Indian Ocean stock (see Rosenbaum *et al.*, 2006b), perhaps as a result of whaling activities during the XIX and XX centuries and/or a lower gene flow with other southern

stocks. The latter explanation is concordant with the lower number of shared haplotypes found in this population respect to other South Pacific stocks (Olavarria *et al.*, 2007).

The results of the AMOVA analysis suggesting panmixia within the breeding grounds of the Breeding Stock G (Ecuador and Colombia) is supported at the demographic level by analyses of photo-identified individuals (Flórez-González *et al.*, 1998; Castro *et al.*, 2008), although a more thorough comparison is needed, including larger sample size catalogues from these breeding areas and others further north (Costa Rica and Panama). However, when the haplotype dataset was stratified by sex and year the AMOVA analysis showed significant differences at haplotype and nucleotide composition, which suggest that the population would not have a complete panmictic distribution. Differences in sampling timing between females in 2006 and 2007 were responsible of such heterogeneity. Heterogeneity within humpback whale Southern Breeding Stocks was reported by Rosenbaum *et al.* (2006) when data were stratified by sex, year and substocks, although contrarily to our findings the authors reported a higher level of structure in males.

While heterogeneity in Ecuadorian whales could be the result of bias caused by a small sample size, variability in the migrating behaviour cannot be excluded. Evidence of such variability exists for both Northern and Southern populations (see Dawbin, 1966; Craig and Herman, 1997). It has been found, for example, that the migration regime in Hawaiian humpback whales is different for each sex, with males undertaking or completing the winter migration more often than females (Craig and Herman, 1997). Our results support this belief, with males arriving to the breeding area off Ecuador in an annual basis and females with a periodicity still to be defined. Even more, females would not be arriving in a random manner but with a year to year variation as showed by the different haplotypes found in each of two compared years. Such level of structure has not been reported in humpback whales before and may be related with a coordinated migration of whales from the same feeding areas. Influence of maternal lineages in feeding aggregations has been found in humpback whales in the North Atlantic (Weinrich *et al.* 1996) and also in the Southeast Pacific at Magellan Strait (Olavarria *et al.*, 2006). It is possible that such influence maintains during the breeding migration and may last up to an early stage at breeding grounds. After their arrival, whales would distribute differently, since no relatedness has been found in breeding groups (Pomilla and Rosenbaum, 2006). Although the breeding migration is in function of the hormonal state, other aspects may influence females' choice to not complete the annual migration or not to undertake it at all, including the energetic costs involved in gestation and lactation and the return to feeding grounds before reaching the breeding area if pregnancy occurs during the migration (Craig and Herman, 1997). The issue requires further research because its potential implications in population assessments and modeling.

Most sampled individuals in Ecuador share haplotypes with whales from Colombia, Magellan Strait and the Antarctic Peninsula. As it was expected, the most common haplotype found in the samples from Ecuador (SP90) is the most common haplotype found in those sampled areas, likewise occurred with the second and third most common haplotypes (SP32 and SP98). These haplotypes have not been recorded in other stocks of the South Pacific basin (see Olavarria *et al.*, 2007) and therefore could be used to characterize this humpback whale stock. Interestingly, Ecuador and Colombia breeding grounds parallel their genetic relationship with the Antarctic Peninsula, confirming this last as the main feeding area of this stock. Ecuadorian whales, however, differentiated from the Magellan Strait, supporting that heterogeneity would occur in the migratory pattern of the Stock G (Olavarria *et al.*, 2006). This is also supported by photo-identified individuals that show a significant higher correspondence between whales feeding at Magellan Strait with whales breeding at Panama than with Ecuador and Colombia (Acevedo *et al.*, 2007).

The presence of haplotype SP16 in the sample from Ecuador, which has been found only in the Western Australian population, is puzzling. It has been demonstrated that extensive movement across humpback whale stocks occur, as shown by the movement of marked whales from Western Australia into the South Pacific (Chittleborough, 1965) and of individually identified whales across the South Pacific (SPWRC, 2006; Robbins *et al.*, 2008; Steel *et al.*, *In prep*). Although the absence of haplotype SP16 in the samples from the Eastern and Central Pacific cannot be ruled out due to sampling, an alternative explanation would be a genetic flow from far distant Western Australian whales. Pomilla and Rosenbaum (2005) demonstrated that humpback whales move between the Indian and Atlantic Ocean basins and therefore such a move could also happen between Ecuador and Western Australia. No matches have been found using photo-ID whales from the Southeast Pacific and the Western Atlantic (Stevick *et al.*, 2004), however, a Southeast Pacific characteristic haplotype has been reported in Brazilian breeding ground (Engel *et al.*, 2008). Collaboration between research groups working with this species in the Southern Hemisphere (e.g. Rosenbaum *et al.*, 2006a) will provide a better understanding of the genetic flow of the species at hemispheric scale.

The sex bias found in this study with males outnumbering females (2.5:1) is similar to that reported in other studies carried out at breeding areas (1.7:1 in the North Atlantic, Palsbøll *et al.* (1997); 1.86:1 in Hawaii, Craig and Herman (1997); 2.4:1 in Eastern Australia, Brown *et al.* (1995); 1.95:1 in South Pacific Olavarria *et al.* (2007)). It has been postulated that the sex bias observed at breeding grounds could be related to migration behavior and/or females heterogeneous distribution (Félix and Haase, 2005), given that such a difference does not occur at feeding grounds (Clapham, *et al.*, 1995), neither in the unique population of the Arabian Sea (Mikhalev, 1997). It is unknown how such a disproportion could affect genetic diversity studies based on mtDNA which is maternally transmitted, although Rosenbaum *et al.* (2006b) did not find a measurable effect regarding sex proportion when examined the population structure of several whale stocks in the Southern Hemisphere. The results of our analysis by sex and haplotype composition support the hypothesis of differences in the migrating behavior in this species as a valid explanation for the skewed proportion toward males in this species found in breeding grounds.

ACKNOWLEDGEMENTS

We would like to thank skippers and tourist operators in Salinas, Ecuador for their assistance and particularly their patience during the sampling. Jéssica Falconí, Andrea Auger, Natalia Botero and Ben Haase, assisted in the field. Jennifer Jackson, Debbie Steel and Gisela Caccone read earlier versions of this manuscript and provided valuable comments. The humpback whale research in Ecuador is supported by the Whale and Dolphin Conservation Society (WDCS). Fieldwork in the Antarctic Peninsula was supported by INACH, the SO-GLOBEC program, IWC and CEQUA. Fieldwork in the Magellan Strait was funded by CEQUA. We thank A. Aguayo-Lobo, J. Acevedo, L. Medrano, D. Thiele, H. Rosenbaum, C. Pomilla, J. Loo and C.S. Baker for key participation on the Antarctic Peninsula and Magellan Strait research projects. Laboratory analysis of those samples was supported by a Marsden Fund grant to C.S. Baker. We thank C.S. Baker for support and guidance.

REFERENCES

- Acevedo-Gutiérrez, A. and Smulter, M.A. 1995. First records of humpback whales including calves at Golfo Dulce and Isla del Coco, Costa Rica, suggesting geographical overlap of northern and southern hemisphere populations. *Marine Mammal Science* 11:554-560.
- Acevedo, J., Rasmussen, K., Félix, F., Castro, C., Llano, M., Secchi, E., Saborío, M.T. Aguayo-Lobo, A., Haase, B., Scheidat, M., Dalla-Rosa, L., Olavarria, C., Forestell, P., Acuña, P., Kaufman, G., and Pastene, L. (2007). Migratory destinations of the humpback whales from Magellan Strait feeding ground, Chile. *Marine Mammal Science* 23(2): 453-463.

- Amos, W., Whitehead, H., Ferrari, M.J., Glockner-Ferrari, D.A., Payne, R. and Gordon, J. 1992. Restrictable DNA from sloughed cetacean skin: its potential for use in population analysis. *Marine Mammal Science* 8:275-283.
- Baker, C. S., Flórez-González, L., Abernethy, B., Rosenbaum, H. C., Slade, R.W., Capella, J., and Bannister, J. L. 1998. Mitochondrial DNA variation and maternal gene flow among humpback whales of the Southern Hemisphere. *Marine Mammal Science* 14(4):721-737.
- Brown, M.R. Corkeron, P.J., Hale, P.T., Schultz, K.W. and Bryden, M.M.1995. Evidence for a sex-segregated migration in the humpback whale (*Megaptera novaeangliae*). *Proceeding of the Royal Society of London, Serie B, Biological Science* 259: 229-234.
- Caballero, S., Hamilton, H., Flórez-González, L., Capella, J., Olavaria, C., Rosenbaum, H.C. and Baker, C.S. 2000. Stock identity and diversity of humpback whale mitochondrial DNA lineages on the Colombian winter breeding grounds. Document SC/52/IA14 submitted to the 52nd Scientific Committee of the International Whaling Commission, Adelaide, Australia, June 2000.
- Caballero, S., Hamilton, H., Jaramillo, H., Capella, J., Flórez-González, L., Olavaria, C., Rosenbaum, H. C., Guhl, F. and Baker, C. S. 2001. Genetic characterization of the Colombian Pacific coast humpback whale population using RAPD and mitochondrial DNA sequences. *Memoirs of the Queensland Museum* 47(2):459-464.
- Capella J., Gibbons, J., Flórez-González, L., LLano, M., Valladares, C., Sabaj, V. and Vilina, Y. 2008. Migratory round-trip of individually identified humpback whales at the strait of Magellan: clues on transit times and phylopatry to destinations. *Revista Chilena de Historia Natural* 81: 547-560
- Castro, C., Acevedo, J., Allen, J.M., Dalla Rosa, L., Flórez-González, L., Aguayo-Lobo, A., Rassmusen, K., Llano, M., Garita, F., Forestell, P., Secchi, E., García-Godos, I., Ferina, D., Kaufman, G., Scheidat, M. and Pasterne, L. 2008. Migratory movements of humpback whales (*Megaptera novaeangliae*) between Machalilla National Park, Ecuador and Southeast Pacific. Document SC/60/SH23, submitted to the Scientific Committee of the International Whaling Commission, Santiago, Chile, 1-13 June 2008. 6pp.
- Chittleborough, R.G. 1965. Dynamics of two populations of humpback whales, *Megaptera novaeangliae* (Borowski). *Australian Journal of Marine and Freshwater Research* 16:33-128.
- Clapham, P.J., Berube, M. and Matilla, D.K. 1995. Sex ratio of the Gulf of Maine humpback whale population. *Marine Mammal Science* 11(2): 227-231.
- Craig, A.S. and Herman, L. 1997. Sex differences in site fidelity and migration of humpback whales (*Megaptera novaeangliae*) to the Hawaiian Islands. *Canadian Journal of Zoology* 75(11): 1923–1933 (1997).
- Engel, M.H., N.J.R. Fagundes, H.C. Rosenbaum, M.S. Leslie, P.H. Ott, R. Schmitt, E. Secchi, L. Dalla Rosa and S. L. Bonatto. 2008. Mitochondrial DNA diversity of the Southwestern Atlantic humpback whale (*Megaptera novaeangliae*) breeding area off Brazil, and the potential connections to Antarctic feeding areas. *Conservation Genetics* 9(5): 1253-1262.
- Excoffier, L. 1995. AMOVA: Analysis of Molecular Variance. University of Geneva, Geneva
- Félix, F. and Haase, B. 2001. The humpback whale off the coast of Ecuador, population parameters and behavior. *Revista de Biología Marina y Oceanografía*, 36(1):61-74.
- Félix, F. and Haase, B. 2005. Distribution of humpback whales along the coast of Ecuador and management implications. *Journal of Cetacean Research and Management*. Vol. 7(1):21-31.
- Félix, F., Palacios, D., Caballero, S., Haase, B. and Falconí, J. 2006a. The 2005 Galápagos humpback whale expedition: a first attempt to assess and characterize the population in the archipelago. Document SC/A06/HW15 submitted to the International Whaling Commission Workshop on Comprehensive Assessment of Southern Hemisphere Humpback Whales, Hobart, Tasmania: 3-7 April 2006. 7p.
- Félix, F., Castro, C., Haase, B., Forestell, P., Álava, J. J. and Scheidat, M. 2006b. Estimates of the Southeastern Pacific humpback whale stock with mark-recapture models in Ecuador. Document SC/A06/HW13 submitted to the International Whaling Commission Workshop on Comprehensive Assessment of Southern Hemisphere Humpback Whales, Hobart, Tasmania: 3-7 April 2006. 7p.
- Félix, F., Caballero S. and Olavaria, C. 2007a. A preliminary assessment of the genetic diversity in humpback whales (*Megaptera novaeangliae*) from Ecuador and population differentiation with other southern hemisphere breeding grounds and feeding areas. Document SC/59/SH11 submitted to the 59th Scientific Committee of the International Whaling Commission, Anchorage, USA, 7-18 May 2007.
- Félix, F., Falconí, J. Auger, A., Haase B. and Jiménez, P. 2007b. Scientific and educational activities in Ecuador during 2006. Final Report to Whale and Dolphin Society WDCS, England. Guayaquil, Ecuador, January 2007. 24 p. (Unpublished).

- Flórez-González, L., Capella, J., Haase, B., Bravo, G. A. Félix, F. and Gerrodette, T. 1998. Changes in winter destinations and the northernmost record of southeastern Pacific humpback whales. *Marine Mammal Science* 14(1):189-196.
- Garrigue, C., Aguayo, A., Amante-Helweg, V.L.U., Baker, C.S., Caballero, S., Clapham, P., Constantine, R., Denkinger, J., Donoghue, M., Flórez-González, L., Greaves, J., Hauser, N., Olavarria, C., Pairoa, C. Peckham, H. and Poole, M. 2002. Movements of humpback whales in Oceania, South Pacific. *Journal of Cetacean Research and Management* 4:255-260.
- Gilson, A., Syvanen, M., Levine, K. and Banks, J. 1998: Deer gender determination by Polymerase Chain Reaction: validation study and application to tissues, bloodstains and hair forensic samples from California. *California Fish and Game*, 84: 159-169.
- IWC. 2001. Report of the Sub-Committee on comprehensive assessment of Southern humpback whales. Report of the Scientific Committee. Annex G. *Reports of the International Whaling Commission*. 48:170-182.
- Katona, S., Baxter, B., Brazier, O., Kraus, S., Perkins, J. and Whitehead, S. 1979. Identification of humpback whales by fluke photographs. Pages 33-44 in H.E. Winn and B.L. Olla, eds. Behavior of Marine Animals. Plenum Press, New York.
- Kellogg, R. 1929. What is known of the migration of some of the whalebone whales. *Smithsonian Institution Annual Report* 1928:467-494.
- Lambertsen, R.H. 1987. A biopsy system for large whales and its use for cytogenetics. *Journal of Mammalogy* 68:443-445.
- Mackintosh, N.A. 1942. The southern stocks of whalebone whales. *Discovery Reports* 22:197-300.
- Maddison, D.R. and Maddison, W.P. (2000) *MacClade: analysis of phylogeny and character evolution*. Sinauer, Sunderland, MA, USA.
- Mikhalev, Y.A. 1997. Humpback whales *Megaptera novaeangliae* in the Arabian Sea. *Marine Ecology Progress Series*, 149:13-21.
- Olavarria C., Baker C.S., Medrano, L., Aguayo-Lobo, A., Caballero, S., Flórez-González, L., Capella, J., Rosenbaum, H.C., Garrigue, C. and Graves, J. 2000. Stock identity of Antarctic Peninsula humpback whales inferred from mtDNA variation. Document SC/52/IA15 submitted to the Scientific Committee of the International Whaling Commission. Adelaide, Australia, June 2000.
- Olavarria, C., Aguayo, A., Acevedo, J., Medrano, L., Thiele, D. and Baker, C.S. 2006. Genetic differentiation between two feeding areas of the Eastern South Pacific humpback whale population: Update on SC/57/SH3. Document SC/A06/HW29 submitted to the International Whaling Commission Workshop on Comprehensive Assessment of Southern Hemisphere Humpback Whales, Hobart, Tasmania: 3-7 April 2006. 7p.
- Olavarria, C., Baker, C.S., Garrigue, C., Poole, M., Hauser, N., Caballero, S., Flórez-González, L., Brasseur, M., Bannister, J., Capella, J., Clapham, P., Dodemont, R., Donoghue, M., Jenner, C., Jenner, M. N., Moro, D., Oremus, M., Paton, D., Rosenbaum, H. and Russell, K. 2007. Population structure of South Pacific humpback and the origin of the eastern Polynesian breeding grounds. *Marine Ecology Progress Series* 330:257-268.
- Omura, H. 1953. Biological study on humpback whales in the Antarctic whaling areas IV and V. *Scientific Report of the Whales Research Institute* 8:81-102.
- Palsbøll, P.J., Allen, J., Bérubé, M., Clapham, P. J., Feddersen, T.P., Hammond, P.S., Hudson, R.R., Jørgensen, H., Katona, S., Larsen A.H., Larsen, F., Lien, J., Matilla, D.K., Sigurjónsson, J., Sears, R., Smith, T., Sponer, R., Stevick, P. and Øien, N. 1997. Genetic tagging of humpback whales. *Nature*, 388:767-769.
- Palumbi, S. R. 1996. Nucleic acids II: the polymerase chain reaction. In: Molecular Systematics (eds, Hillis D, Moritz C, Mable BK) Sinauer Associates, Sunderland, MA.
- Papastavrou, V. and Van Waerebeek, K. 1997. A note on the occurrence of humpback whales (*Megaptera novaeangliae*) in tropical and subtropical areas: the upwelling link. *Reports of the International Whaling Commission* 47:945-947
- Pomilla, C. and Rosenbaum, H. 2005. Against the current: an inter-oceanic whale migration event. *Biology Letters* 1:476-479.
- Pomilla, C. and Rosenbaum, H. 2006. Estimates of relatedness in groups of humpback whales (*Megaptera novaeangliae*) on two wintering grounds of the Southern Hemisphere. *Molecular Ecology* 15:2541-2555.
- Ramírez, P. 1988. La ballena jorobada *Megaptera novaeangliae* en la costa norte del Perú: Períodos 1961-1965 y 1975-1985. *Boletín de Lima* 56:91-96. (In Spanish).
- Rasmussen, K., Palacios, D.M., Calambokidis, J., Saborio, M.T., Dalla Rosa, L., Secchi, E.R., Steiger, G.H., Allen, J.M. and Stone, G.S. 2007. Southern Hemisphere humpback whales wintering off

- Central America: insights from water temperature into the longest mammalian migration. *Biology Letters* 3(3): 302-305.
- Raymond, M. and Rousset, F. 1995. An exact test for population differentiation. *Evolution* 49:1280-1283.
- Robbins, J., Dalla Rosa, L., Allen, J.M., Mattila, D.K. and Secchi, E.R. 2008. Humpback whale photo-identification reveals exchange between American Samoa and the Antarctic Peninsula, and a new mammalian distance record. Document SC/60/SH5, submitted to Scientific Committee of the International Whaling Commission, Santiago, Chile, 1-13 June 2008.
- Rosenbaum, H.C., Pomilla, C., Olavarria, C., Baker, C.S., Leslie, M.C., Mendez, M.C., Caballero, S., Brassuer, M., Bannister, J., Best, P.B., Bonatto, S., Collins, T., Engel, M.H., Ersts, P.J., Findlay, K.P., Florez-Gonzalez, L., Garrigue, C., Hauser, N., Jenner, C., Meyer, M., Minton, G., Poole, M. and Razafindrakoto, Y. 2006a. A first and preliminary analysis of mtDNA sequences from humpback whales for breeding stocks A-G and X. Document SC/A06/HW59 submitted to the Inter-session workshop for the Comprehensive Assessment of Southern Hemisphere humpback whales Scientific Committee of the International Whaling Commission. Hobart, 3 - 7 April 2006.
- Rosenbaum, H.C., Pomilla, C., Olavarria, C., Leslie, M.C., Mendez, M.C., Collins, T., Engel, M.H., Ersts, P.J., Findlay, K.P., Bonatto, S., Kotze, P.G.H., Meyer, M., Minton, G., Berendse, J., Thorton, M., Razafindrakoto, Y. and Ngouessono, S. 2006b. Mitochondrial DNA diversity and population structure of humpback whales from their wintering areas (breeding stocks) in the Indian and South Atlantic oceans (wintering regions A,B,C and X). Document SC/A06/HW41 submitted to the Inter-session workshop for the Comprehensive Assessment of Southern Hemisphere humpback whales Scientific Committee of the International Whaling Commission. Hobart, 3 - 7 April 2006.
- Saiki, R.K., Gelfand, D.H., Stofell, S., Scharf, R., Higuchi, R., Horn, G.T., Mullis, K.B. and Erlich, H.A. 1988. Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science*, 239, 487-491.
- Schneider, S., Roessli D. and Excoffier, L. 2006. Arlequin Ver 3.1, An integrate software package for population Genetics. <http://cmpg.unibe.ch/software/arlequin3>.
- SPWRC, Garrigue, C., Baker, C.S., Constantine, R., Poole, M., Hauser, N., Clapham, P., Donoghue, M., Russell, K., Paton, D. and Mattila, D. 2006. Interchange of humpback whales in Oceania (South Pacific), 1999 to 2004. Document SC/A06/HW55 submitted to the Inter-session workshop for the Comprehensive Assessment of Southern Hemisphere humpback whales Scientific Committee of the International Whaling Commission. Hobart, 3 - 7 April 2006.
- Steel, D., Garrigue, C., Poole, M.M., Hauser, N., Olavarria, C., Flórez-González, L., Constantine, R., Caballero, S., Thiele, D., Slooten, L., Dawson, S., Oremus, M., Russell, K., Paton, D., Clapham, P., Donoghue, M. and Baker, C.S. *In preparation*. Migratory connections between humpback whales from South Pacific breeding grounds and Antarctic feeding areas based on genotype matching.
- Stevick, P., Aguayo, A., Allen, J., Avila, I. C., Capella, J., Castro, C., Chater, K., Engel, M. H., Félix, F., Flórez-González, L., Freitas, A., Haase, B., Llano, M., Lodi, L., Muñoz, E., Olavarria, C., Secchi, E., Scheidat, M. and Siciliano, S. 2004. A note on the migrations of individually identified humpback whales between the Antarctic Peninsula and South America. *Journal of Cetacean Research and Management*, 6(2):109-113.
- Stone, G., Flórez-González, L. and Katona, S. 1990. Whale migration record. *Nature* 346: 705.
- Townsend, C.H. 1935. The distribution of certain whales as shown by logbook records of American whaleships. *Zoologica NY* 16:133-144.
- Weinrich, M. T., Rosenbaum, H., Baker, C.S., Blackmer, A.L. and Whitehead, H. 2006. The influence of maternal lineages on social affiliations among humpback whales (*Megaptera novaeangliae*) on their feeding grounds in the Southern Gulf of Maine. *Journal of Heredity* 97(3):226-234.